

# Radio detection of cosmic rays with LOFAR

Stijn Buitink

- Radio emission from air showers
- LOFAR analysis pipeline
- First analysis results

A. Corstanje, J.E. Enriquez, H. Falcke, W. Frieswijk, J. Hörandel, T. Huege, M. Krause, A. Nelles, P. Schellart, O. Scholten, S. Thoudam, S. ter Veen, M. van den Akker  
for the **LOFAR** collaboration



LOFAR

Radboud Universiteit Nijmegen





university of  
 groningen





# Radio: a new handle on CR Composition




Auger ICRC 2013 Rio de Janeiro

**Surface detection:** muon component  
- very sensitive to hadronic interaction models

**Fluorescence Detection** to measure  $X_{\max}$   
- duty cycle  $\sim 10\%$ ,  $\sigma \sim 20$  g/cm<sup>2</sup>


**Radio Detection** to measure  $X_{\max}$ :  
- cheap detectors, duty cycle  $\sim 100\%$   
- attractive method to increase statistics at ultra-high energy!



*Does it work? Is the precision good enough?*

# What drives the radio emission?

- Earth magnetic field  
electrons/positrons deflected  
 $E \sim dn_{ch}/dt$
- Charge excess  
negative charge due to electron knockouts  
 $E \sim d(n_e - n_p)/dt$
- Non-unity index of refraction  
Cherenkov-like effects  
ring structure possible



Coherent at 100 MHz  
wavelength  $>$  shower front size  
 $P \sim n^2$






# LOFAR

low frequency array  
10 - 250 MHz

Epoch of Reionization  
Radio Transients  
Astroparticle Physics  
Cosmic Magnetism  
Surveys  
Solar Physics





**SUPERTERP**  
 ~600 low band antennas  
 10 - 80 MHz  
 5 ns time resolution  
 > GB buffer/antenna

+ LORA

LOFAR Radboud air shower array  
 20 scintillator stations (ex-KASCADE)

24 core stations  
 9 remote stations  
 8 international stations




# Reconstruction pipeline







1. LORA scintillators provide trigger & shower reco



cleaning  
→




2. RFI cleaning using phase stability of emitters




# Particle Detector & Radio Agree

P. Schellart




3. pulse finding

Curvature



4. arrival direction  
fit




Geomagnetic  
= dominant

5. Apply antenna model to get LDF




# Air Showers at 30 - 80 MHz



Scintillator threshold

range  $10^{16.5} - 10^{18.5}$  eV  
north-south asymmetry



sample of 400+ showers




**Xmax measurement by fitting  
lateral distribution of P**



# radio pattern



vector sum of **geomagnetic** and **charge excess** component  
relativistic beaming  
distortion by Cherenkov-like effects ( $n \neq 1$ )







CoREAS simulation



$X_{\max} \sim 600 \text{ g/cm}^2$

$X_{\max} \sim 650 \text{ g/cm}^2$

$X_{\max} \sim 700 \text{ g/cm}^2$




LOFAR:  
200 - 400 antennas/event

→ *fit full 2D pattern !*



# Strategy



# Fit for each simulation

Minimize  $\chi^2$  of **radio** and **particle** data simultaneously

$$\chi^2 = \sum_{\text{antennas}} \left( \frac{P_{\text{ant}} - f_r P_{\text{sim}}(x_{\text{ant}} + x_{\text{off}}, y_{\text{ant}} + y_{\text{off}})}{\sigma_{\text{ant}}} \right)^2 + \sum_{\text{detectors}} \left( \frac{d_{\text{det}} - f_p d_{\text{sim}}(x_{\text{det}} + x_{\text{off}}, y_{\text{det}} + y_{\text{off}})}{\sigma_{\text{det}}} \right)^2$$

4 fit parameters:


core position

radio power scale factor

particle density scale factor




# best fit out of 40 simulations




# best fit out of 40 simulations

Lateral distribution radio signal



1D LDFs don't fit !


LORA ldf after core optimization



- core shifted by ~ 50m
- revised energy estimate



Fit Quality after core optimization



# Errors

## Procedure:

- single out 1 sim
  - add noise
  - reconstruct with other 39 sims
  - calculate error  $\Delta X$
- repeat for all sims


construct region that contains 68% of  $\Delta X$

$$\sigma_{\text{meth}} = 12.7 \text{ g/cm}^2$$


$$\sigma_{\text{atm}} = 15 \text{ g/cm}^2$$

$$\sigma = 19.6 \text{ g/cm}^2$$


other systematic effects under investigation




Radiation profile in shower plane






Lateral distribution radio signal



Fit Quality after core optimization







# Conclusions



- **We understand air shower radio emission**
  - CoREAS sim in agreement with 300+ data point events!
  - geomagnetic, charge excess & Cherenkov observed
  - absolute power not yet available...
- **We can measure  $X_{\max}$** 
  - accurate reconstruction based on CoREAS sim  $\sigma \sim 20 \text{ g/cm}^2$
  - *coming*: independent analysis using wavefront shape
  - radio has duty cycle of near 100%
- **Future**
  - LOFAR composition energy range  $10^{16.5} - 10^{18.5} \text{ eV}$
  - TUNKA-REX comparison with air-Cherenkov
  - AERA (Auger) comparison with FD & composition at highest energies

research funded by ERC Advanced Grant (H. Falcke) & NWO Veni Grant (S. Buitink)

# backup slides



# polarization





Total power

X-dipole


Y-dipole

# delays w.r.t. plane wave

- Wavefront curvature  
nanosecond precision  
(sub-sample!)
- not purely spherical or  
conical
- sensitive to  $X_{\max}$   
(analysis ongoing)



# Energy Calibration




- signal strength at 60-70m axis distance
- coherent signal  $P \sim n^2$
- spread due to shower-to-shower fluctuations



# radio emission: converging models

microscopic:



geosynchrotron +  
'endpoint' radiation



macroscopic:

time-dependent  
current densities

$$E \sim dJ/dt \sim dN/dt$$






Huege, Ludwig Scholten, De Vries NIMPA 662 (2012) 179

Now available: CoREAS, ZHAireS, EVA, Selfas  
all include geomagnetic, charge excess, cherenkov effects

# simulations

- CORSIKA 6.990 QGSJETII-03 + FLUKA
- CoREAS v0.9 T. Huege et al. AIPC 1535, 128 (2013)



raw sim pulses  
(xyz)

antenna simulation  
(incl. 30-80 MHz filt.)

power in  
X/Y dipole